

RECEIVED
CENTRAL FAX CENTER

JUL 27 2006

Atty Dkt. No.: 10030416-1
USSN: 10/698,195

AMENDMENTS TO THE CLAIMS

Please incorporate the following amendments to the subject application.

In the Claims:

1. **(Currently Amended)** A method of identifying a sequence of a nucleic acid for use as a substrate surface immobilized probe for a target nucleic acid, said method comprising:
 - (a) determining via an algorithm a full length synthesis probability measure for each member sequence of a set of a plurality of candidate probe sequences for said target nucleic acid; and
 - (b) employing said determined full length synthesis probability measures to identify and select a sequence for use as a substrate immobilized probe for said target nucleic acid.
2. **(Original)** The method according to Claim 1, wherein those sequences of said set that satisfy a full length synthesis probability threshold are selected .
3. **(Original)** The method according to Claim 2, wherein said full length synthesis probability measure is a measure of the probability that said candidate probe sequence will be fully synthesized by an in situ nucleic acid synthesis protocol.
4. **(Original)** The method according to Claim 3, wherein said full length synthesis probability measure is an evaluation of said candidate probe sequence's depurination susceptibility during said in situ nucleic acid synthesis protocol.
5. **(Original)** The method according to Claim 4, wherein said depurination susceptibility measure is a deblock dose or parameter based thereon.
6. **(Original)** The method according to Claim 5, wherein said method comprises determining a total deblock dose for each member sequence of said set.

Atty Dkt. No.: 10030416-1
USSN: 10/698,195

7. (Original) The method according to Claim 6, wherein said full length synthesis probability threshold is a total deblock dose threshold and selected sequences have a total deblock dose that does not exceed said deblock dose threshold.

8. (Original) The method according to Claim 6, wherein said total deblock dose is a sum of individual deblock doses over all A nucleotides of said candidate probe sequence, except for any 5' terminal A nucleotide.

9. (Original) The method according to Claim 8, wherein each of said individual deblock doses $d(x)$ for an A nucleotide positioned at layer x of a candidate probe sequence to be produced by a process having L total layers is equal to $L-x+1$.

10. (Original) The method according to Claim 1, wherein at least some of said steps are carried out by a computational analysis system.

Claims 11 and 12 (Cancelled).

13. (Original) A method of producing a nucleic acid array, said method comprising:
producing at least two different probe nucleic acids immobilized on a surface of a solid support, wherein at least one of said at least two different probe nucleic acids has a sequence of nucleotides identified according to the method of Claim 1.

14. (Original) The method according to Claim 13, wherein said at least two different probe nucleic acids are produced on said surface of said solid support by synthesizing said probe nucleic acids on said surface.

15. (Original) The method according to Claim 13, wherein said at least two different probe nucleic acids are produced on said surface of said solid support by depositing said at least two different probe nucleic acids onto said surface of said solid support.

Atty Dkt. No.: 10030416-1
USSN: 10/698,195

Claims 16-21 (Cancelled).

22. (Original) In a method of identifying a sequence of a nucleic acid that is suitable for use as a substrate surface immobilized probe for a target nucleic acid, the improvement comprising:

evaluating a plurality of candidate probe sequences for said target nucleic acid for depurination susceptibility.

23. (Cancelled).

24. (New) A method of identifying a sequence of a nucleic acid for use as a substrate surface immobilized probe for a target nucleic acid, said method comprising:

(a) determining a full length synthesis probability measure for each member sequence of a set of a plurality of candidate probe sequences for said target nucleic acid by evaluation of the susceptibility to depurination during synthesis of each probe sequence; and

(b) employing said determined full length synthesis probability measures to identify and select a sequence for use as a substrate immobilized probe for said target nucleic acid.

25. (New) The method according to Claim 24, wherein said depurination susceptibility measure is a deblock dose or parameter based thereon.

26. (New) The method according to Claim 25, wherein said method comprises determining a total deblock dose for each member sequence of said set.

27. (New) The method according to Claim 26, wherein said full length synthesis probability threshold is a total deblock dose threshold and selected sequences have a total deblock dose that does not exceed said deblock dose threshold.

Atty Dkt. No.: 10030416-1
USSN: 10/698,195

28. **(New)** The method according to Claim 26, wherein said total deblock dose is a sum of individual deblock doses over all A nucleotides of said candidate probe sequence, except for any 5' terminal A nucleotide.

29. **(New)** The method according to Claim 28, wherein each of said individual deblock doses $d(x)$ for an A nucleotide positioned at layer x of a candidate probe sequence to be produced by a process having L total layers is equal to $L-x+1$.